

346



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : A01N 43/40, A61K 31/445</p>	<p>A1</p>	<p>(11) International Publication Number: WO 98/31225 (43) International Publication Date: 23 July 1998 (23.07.98)</p>
<p>(21) International Application Number: PCT/US98/00618 (22) International Filing Date: 13 January 1998 (13.01.98) (30) Priority Data: 60/035,591 17 January 1997 (17.01.97) US (71) Applicant: BRISTOL-MYERS SQUIBB COMPANY [US/US]; P.O. Box 4000, Princeton, NJ 08543-4000 (US). (72) Inventor: GREGG, Richard, E.; 7 Linden Lane, Pennington, NJ 08534 (US). (74) Agents: RODNEY, Burton et al.; Bristol-Myers Squibb Com- pany, P.O. Box 4000, Princeton, NJ 08543-4000 (US).</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report.</i></p>
<p>(54) Title: A METHOD OF INHIBITING OR TREATING PHYTOSTEROLEMIA WITH AN MTP INHIBITOR (57) Abstract A method is provided for inhibiting onset of or treating phytosterolemia by administering to a patient an MTP inhibitor, alone or optionally, in combination with another cholesterol lowering drug, such as pravastatin.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

A METHOD OF INHIBITING OR TREATING PHYTOSTEROLEMIA WITH AN MTP INHIBITOR

5 Field of the Invention

The present invention related to a method for inhibiting onset of or treating phytosterolemia, by administering an MTP inhibitor alone or in combination with another cholesterol lowering drug, such as pravastatin.

10

Background of the Invention

As indicated in Scriver, C.R. et al "Metabolic Basis of Inherited Diseases" Vol. II (1995), Chap. 65, Inborn Errors in Bile Acid Biosynthesis and Storage of Sterols
15 Other than Cholesterol, Bjorkhem, I. and Boberg, K.M., pp. 2073-2099, phytosterolemia (also referred to as sitosterolemia) is a rare inherited sterol storage disease involving increased intestinal absorption of phytosterol or shellfish sterols and decreased fecal secretion. It is
20 characterized by "tendon and tuberous xanthomas and by a strong predisposition to premature coronary atherosclerosis.... Increased amounts of phytosterols (plant sterols), such as sitosterol and campesterol and their 5 α -stanols, are found in blood, plasma, erythrocytes,
25 and different tissues, especially in the xanthomas and arteries of affected subjects. Increased serum cholesterol and cholesterol have also been found in many patients."
(p. 2073)

Patients afflicted with phytosterolemia have been
30 found to have an increased incidence of coronary heart disease at an early age most likely due to early development of atherosclerosis at an early age. Bjorkhem et al, supra, indicate at page 2090 that "the mechanism behind the atherosclerosis is unexplained, but a high
35 content of plant sterols in the circulating lipoproteins might promote their deposition in the arterial wall."

The microsomal triglyceride transfer protein (MTP) catalyzes the transport of triglyceride (TG), cholesteryl ester (CE), and phosphatidylcholine (PC) between small unilamellar vesicles (SUV). Wetterau & Zilversmit, Chem. Phys. Lipids 38, 205-22 (1985). When transfer rates are expressed as the percent of the donor lipid transferred per time, MTP expresses a distinct preference for neutral lipid transport (TG and CE), relative to phospholipid transport. The microsomal triglyceride transfer protein from bovine liver has been isolated and extensively characterized (1). This has led to the cloning of cDNA expressing the protein from several species, including humans (2). MTP is composed of two subunits. The small subunit is the previously characterized multifunctional protein, protein disulfide isomerase. This is supported by biochemical analysis of the protein (3) as well as co-expression studies performed in insect Sf9 cells using the baculovirus expression system. Expression of soluble active MTP requires the co-expression of PDI and the unique large subunit of MTP (4).

1: Wetterau, J.R. and Zilversmit, D.B. (1985) Chem. Phys. Lipids 38, 205-222.

Wetterau, J.R., et al, (1990) J. Biol. Chem. 265, 9800-9807.

Wetterau, J.R., et al, (1991) Biochemistry 30, 4406-4412.

Atzel, A., and Wetterau, J.R. (1993) Biochemistry 32, 10444-10450.

Atzel, A., and Wetterau, J.R. (1994) Biochemistry 33, 15382-15388.

Jamil, H., et al, (1995) J. Biol. Chem. 270, 6549-6554.

2. Sharp, D. et al, (1993) Nature 365, 65-69.

Lin, M.C.M., et al, J. Biol. Chem. 269, 29138-29145.

Nakamuta, M., et al, (1996) Genomics 33, 313-316.

3. Wetterau, J.R., et al, (1990) J. Biol. Chem. 265, 9800-9807.

5 Wetterau, J.R., et al, (1991) Biochemistry 30, 9728-9735.

4. Ricci, B., et al, (1995) J. Biol. Chem. 270, 14281-14285.

10 In vitro, MTP catalyzes the transport of lipid molecules between phospholipid membranes. Presumably, it plays a similar role in vivo, and thus plays some role in lipid metabolism. The subcellular (lumen of the microsomal fraction) and tissue distribution (liver and intestine) of
15 MTP have led to speculation that it plays a role in the assembly of plasma lipoproteins, as these are the sites of plasma lipoprotein assembly. Wetterau & Zilversmit, Biochem. Biophys. Acta 875, 610-7 (1986). The ability of MTP to catalyze the transport of TG between membranes is
20 consistent with this hypothesis, and suggests that MTP may catalyze the transport of TG from its site of synthesis in the endoplasmic reticulum (ER) membrane to nascent lipoprotein particles within the lumen of the ER.

 Abetalipoproteinemia is an autosomal recessive
25 disease characterized by a virtual absence of plasma lipoproteins which contain apolipoprotein B (apoB). Kane & Havel in The Metabolic Basis of Inherited Disease, Sixth edition, 1139-64 (1989). Plasma TG levels may be as low as a few mg/dL, and they fail to rise after fat ingestion.
30 Plasma cholesterol levels are often only 20-45 mg/dL. These abnormalities are the result of a genetic defect in the assembly and/or secretion of very low density lipoproteins (VLDL) in the liver and chylomicrons in the intestine. The molecular basis for this defect had not
35 been previously determined. In subjects examined, triglyceride, phospholipid, and cholesterol synthesis appear normal. At autopsy, subjects are free of

atherosclerosis. Schaefer et al., Clin. Chem. 34, B9-12 (1988). A link between the apoB gene and abetalipoproteinemia has been excluded in several families. Talmud et al., J. Clin. Invest. 82, 1803-6 (1988) and Huang
5 et al., Am. J. Hum. Genet. 46, 1141-8 (1990).

Recent reports (5) demonstrate that the defect causing abetalipoproteinemia is in the MTP gene, and as a result, the MTP protein. When examined, individuals with abetalipoproteinemia have no MTP activity, as a result of
10 mutations in the MTP gene, some of which have been characterized. These results indicate that MTP is required for the synthesis of apoB containing lipoproteins, such as VLDL, the precursor to LDL. It therefore follows that inhibitors of MTP would inhibit the synthesis of VLDL and
15 LDL, thereby lowering VLDL levels, LDL levels, cholesterol levels, and triglyceride levels in animals and man.

5. Wetterau, J.R., et al, (1992) Science 258, 999-1001.
Sharp, D., et al, (1993) Nature 365, 65-69.
20 Ricci, B., et al, (1995) J. Biol. Chem. 270, 14281-14285.

Shoulders, C.C., et al, (1993) Hum. Mol. Genetics 2, 2109-2116.

Narcisi, T.M.E., et al, (1995) Am. J. Hum. Genet.
25 57, 1298-1310.

Rehberg, E.F., et al, J. Biol. Chem (in press).

Canadian Patent Application No. 2,091,102 published March 2, 1994 (corresponding to U.S. application Serial No.
30 117,362, filed September 3, 1993 (file DC21b)) which is incorporated herein by reference), reports MTP inhibitors which also block apoB containing lipoprotein secretion in a human hepatic cell line (HepG2 cells). This provides further support for the proposal that an MTP inhibitor
35 would lower apoB containing lipoprotein and lipid levels in vivo. This Canadian patent application discloses a method for identifying the MTP inhibitors.

The use of microsomal triglyceride transfer protein (MTP) inhibitors for decreasing serum lipids including cholesterol and triglycerides and their use in treating atherosclerosis, obesity, hyperglycemia, and pancreatitis is disclosed in WO 96/26205, published August 29, 1996, U.S. Application Serial No. 472,067, filed June 6, 1995 (file DC21e), U.S. Application Serial No. 548,811, filed January 11, 1996 (file DC21h), U.S. provisional application No. 60/017,224, filed May 9, 1996 (file HX79a*), U.S. provisional application No. 60/017,253, filed May 10, 1996 (file HX82*), U.S. provisional application No. 60/017,254, May 10, 1996 (file HX84*) and U.S. provisional application No. 60/028,216, filed October 1, 1996 (file HX86*).

All of the above U.S. applications are incorporated herein by reference.

Description of the Invention

In accordance with the present invention, a method is provided for inhibiting onset of or treating phytosterolemia, in mammalian species, wherein a therapeutically effective amount of a microsomal triglyceride transfer protein (MTP) inhibitor is administered to a patient in need of treatment.

The MTP inhibitor may optionally be administered in combination with another cholesterol lowering drug or delipidating agent.

The MTP inhibitor alone or optionally in combination with another cholesterol lowering drug is administered systemically, such as orally or parenterally or transdermally, to patients in need of treatment.

In accordance with the present invention, the MTP inhibitor lowers plasma cholesterol (LDL-cholesterol) to at least about 50% of normal LDL blood level, preferably down to less than about 25% of normal, and lowers triglycerides to at least about 50% of normal triglyceride blood level, and preferably down to about 25% or less of normal, and


thereby reduces plasma cholesterol and resulting atherosclerosis.

The terms "another cholesterol lowering drug or agent" or "another delipidating drug" will be employed interchangeably herein.


MTP inhibitors to be employed in the methods of the invention include MTP inhibitors disclosed in Canadian Patent Application No. 2,091,102 described hereinbefore (corresponding to U.S. Application Serial No. 117,362), WO 92/26205 published August 29, 1996, U.S. Application Serial No. 472,067, filed June 6, 1995 (file DC21e), U.S. Application Serial No. 548,811, filed January 11, 1996 (file DC21h), U.S. provisional application No. 60/017,224, filed May 9, 1996 (file HX79a*), U.S. provisional application No. 60/017,253, filed May 10, 1996 (file HX82*), U.S. provisional application No. 60/017,254, filed May 10, 1996 (file HX84*), and U.S. provisional application No. 60/028,216, filed October 1, 1996 (file HX86*). Preferred are each of the preferred MTP inhibitors disclosed in each of the above applications.

All of the above U.S. applications are incorporated herein by reference.


The MTP inhibitors disclosed in U.S. Application Serial No. 472,067, filed June 6, 1995 (file DC21e) are piperidine compounds of the structure



or




or




30

or



or



where Q is $\text{—}\overset{\text{O}}{\underset{\text{O}}{\text{C}}}\text{—}$ or $\text{—}\overset{\text{O}}{\underset{\text{O}}{\text{S}}}\text{—}$;

5

X is: CHR^8 , $\text{—}\overset{\text{O}}{\underset{\text{O}}{\text{C}}}\text{—}$, $\text{—}\underset{\text{R}^9}{\text{CH}}\text{—}\underset{\text{R}^{10}}{\text{CH}}\text{—}$ or $\text{—}\underset{\text{R}^9}{\text{C}}=\underset{\text{R}^{10}}{\text{C}}\text{—}$;

R^8 , R^9 and R^{10} are independently hydrogen, alkyl, alkenyl, alkynyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl, or cycloalkylalkyl;

10



Y is $\text{—}(\text{CH}_2)_m\text{—}$ or $\text{—}\overset{\text{O}}{\underset{\text{O}}{\text{C}}}\text{—}$

wherein m is 2 or 3;

R^1 is alkyl, alkenyl, alkynyl, aryl, heteroaryl, arylalkyl wherein alkyl has at least 2 carbons, diarylalkyl, arylalkenyl, diarylalkenyl, arylalkynyl, diarylalkynyl, diarylalkylaryl, heteroarylalkyl wherein alkyl has at least 2 carbons, cycloalkyl, or cycloalkylalkyl wherein alkyl has at least 2 carbons, all optionally substituted through available carbon atoms with 1, 2, 3 or 4 groups selected from halo, haloalkyl, alkyl, alkenyl, alkoxy, aryloxy, aryl, arylalkyl, alkylmercapto, arylmercapto, cycloalkyl, cycloalkylalkyl, heteroaryl, fluorenyl, heteroarylalkyl, hydroxy or oxo;


or R^1 is a fluorenyl-type group of the structure

25




5

R^1 is an indenyl-type group of the structure




(a = 2,3 or 4)



10

Z^1 and Z^2 are the same or different and are independently a bond, O, S,




with the proviso that with respect to B, at least one of Z¹ and Z² will be other than a bond; R¹¹ is a bond, alkylene, alkenylene or alkynylene of up to 10 carbon atoms; arylene or mixed arylene-alkylene; R¹² is hydrogen, alkyl, alkenyl, aryl, haloalkyl, trihaloalkyl, trihaloalkylalkyl, heteroaryl, heteroarylalkyl, arylalkyl, arylalkenyl, cycloalkyl, aryloxy, alkoxy, arylalkoxy or cycloalkylalkyl, with the provisos that preferably

- (1) when R¹² is H, aryloxy, alkoxy or arylalkoxy,
 10 then Z² is $\text{—NH—}\overset{\text{O}}{\underset{\text{O}}{\text{C}}}\text{—}$, $\text{—}\overset{\text{O}}{\underset{\text{alkyl}}{\text{N}}}\text{—}\overset{\text{O}}{\underset{\text{O}}{\text{C}}}\text{—}$, $\text{—}\overset{\text{O}}{\underset{\text{O}}{\text{C}}}\text{—}$ or a bond and
 (2) when Z² is a bond, R¹² cannot be heteroaryl or heteroarylalkyl;

Z is a bond, O, S, N-alkyl, N-aryl, or alkylene or alkenylene from 1 to 5 carbon atoms; R¹³, R¹⁴, R¹⁵, and R¹⁶
 15 are independently hydrogen, alkyl, halo, haloalkyl, aryl, cycloalkyl, cycloheteroalkyl, alkenyl, alkynyl, hydroxy, alkoxy, nitro, amino, thio, alkylsulfonyl, arylsulfonyl, alkylthio, arylthio, aminocarbonyl, alkylcarbonyloxy, arylcarbonylamino, alkylcarbonylamino, arylalkyl,
 20 heteroaryl, heteroarylalkyl or aryloxy;

R^{15a} and R^{16a} are independently hydrogen, alkyl, halo, haloalkyl, aryl, cycloalkyl, cycloheteroalkyl, alkenyl, alkynyl, alkoxy, alkylsulfonyl, arylsulfonyl, alkylthio, arylthio, aminocarbonyl, alkylcarbonyloxy,
 25 arylcarbonylamino, alkylcarbonylamino, arylalkyl, heteroaryl, heteroarylalkyl, or aryloxy;


or R¹ is a group of the structure



30 wherein p is 1 to 8 and R¹⁷ and R¹⁸ are each independently H, alkyl, alkenyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl or cycloalkylalkyl at least one of R¹⁷ and R¹⁸ being other than H;

or R¹ is a group of the structure

35



wherein R¹⁹ is aryl or heteroaryl;

R²⁰ is aryl or heteroaryl;

R²¹ is H, alkyl, aryl, alkylaryl, arylalkyl,
 5 aryloxy, arylalkoxy, heteroaryl, heteroarylalkyl,
 heteroarylalkoxy, cycloalkyl, cycloalkylalkyl or
 cycloalkylalkoxy;

R², R³, R⁴ are independently hydrogen, halo, alkyl,
 alkenyl, alkoxy, aryloxy, aryl, arylalkyl, alkylmercapto,
 10 arylmercapto, cycloalkyl, cycloalkylalkyl, heteroaryl,
 heteroarylalkyl, hydroxy or haloalkyl;


R⁵ is independently alkyl, alkenyl, alkynyl, aryl,
 alkoxy, aryloxy, arylalkoxy, heteroaryl, arylalkyl,
 heteroarylalkyl, cycloalkyl, cycloalkylalkyl,
 15 polycycloalkyl, polycycloalkylalkyl, cycloalkenyl,
 cycloheteroalkyl, heteroaryloxy, cycloalkenylalkyl,
 polycycloalkenyl, polycycloalkenylalkyl,
 heteroarylcarbonyl, amino, alkylamino, arylamino,
 heteroarylamino, cycloalkyloxy, cycloalkylamino, all
 20 optionally substituted through available carbon atoms with
 1, 2, 3 or 4 groups selected from hydrogen, halo, alkyl,
 haloalkyl, alkoxy, haloalkoxy, alkenyl, alkynyl,
 cycloalkyl, cycloalkylalkyl, cycloheteroalkyl,
 cycloheteroalkylalkyl, aryl, heteroaryl, arylalkyl,
 25 arylcycloalkyl, arylalkenyl, arylalkynyl, aryloxy,
 aryloxyalkyl, arylalkoxy, arylazo, heteroaryloxo,
 heteroarylalkyl, heteroarylalkenyl, heteroaryloxy, hydroxy,
 nitro, cyano, amino, substituted amino, thiol, alkylthio,
 arylthio, heteroarylthio, arylthioalkyl, alkylcarbonyl,
 30 arylcarbonyl, arylaminocarbonyl, alkoxycarbonyl,
 aminocarbonyl, alkynylaminocarbonyl, alkylaminocarbonyl,
 alkenylaminocarbonyl, alkylcarbonyloxy, arylcarbonyloxy,
 alkylcarbonylamino, arylcarbonylamino, arylsulfinyl,
 arylsulfinylalkyl, arylsulfonyl, alkylsulfonyl,
 35 arylsulfonylamino, heteroarylcarbonylamino,

heteroarylsulfinyl, heteroarylthio, heteroarylsulfonyl, alkylsulfinyl;


R^6 is hydrogen or C_1 - C_4 alkyl or C_1 - C_4 alkenyl; all optionally substituted with 1, 2, 3 or 4 groups which may independently be any of the substituents listed in the definition of R^5 set out above;

R^7 is alkyl, aryl or arylalkyl wherein alkyl by itself or as part of arylalkyl is optionally substituted with oxo $\left(\begin{smallmatrix} \text{O} \\ || \end{smallmatrix} \right)$;

10



are the same or different and are independently selected from heteroaryl containing 5- or 6-ring members; and




15

pharmaceutically acceptable salts thereof; with the provisos that preferably where in the first formula X is CH_2 , and R^2 , R^3 and R^4 are each H, then R^1 will be other than 3,3-diphenylpropyl, and preferably in the fifth formula, where one of R^2 , R^3 and R^4 is 6-fluoro, and the others are H, R^7 will be other than 4-(2-methoxyphenyl).

20

The MTP inhibitors disclosed in U.S. application Serial No. 548,811 filed January 11, 1996 (file DC21h), have the structure



25


including the piperidine N-oxide thereof or a pharmaceutically acceptable salt thereof, wherein Z is a bond, O or S;

x^1 and x^2 are independently selected from H or halo;
 x is an integer from 2 to 6;

R^5 is heteroaryl, aryl, heterocycloalkyl or cycloalkyl, each R^5 group being optionally substituted with
 5 1, 2, 3 or 4 substituents which may be the same or different.

The MTP inhibitors disclosed in U.S. provisional application No. 60/017,224, filed May 9, 1996 (file HX79a*) have the structure

10




including pharmaceutically acceptable salts thereof,
 wherein q is 0, 1 or 2;

A is (1) a bond;



15

(2) $-O-$; or



where R^5 is H or lower alkyl or R^5 together with R^2 forms a carbocyclic or heterocyclic ring system containing 4 to 8
 20 members in the ring.


B is a fluorenyl-type group of the structure:



(the above B is also referred to as a fluorenyl-type ring or moiety); or

25

B is an indenyl-type group of the structure




R^x is H, alkyl or aryl;

R^1 is alkyl, alkenyl, alkynyl, alkoxy, (alkyl or aryl)₃Si (where each alkyl or aryl group is independent), cycloalkyl, cycloalkenyl, substituted alkylamino, substituted arylalkylamino, aryl, arylalkyl, arylamino, aryloxy, heteroaryl, heteroarylamino, heteroaryloxy, arylsulfonylamino, heteroarylsulfonylamino, arylthio, arylsulfinyl, arylsulfonyl, alkylthio, alkylsulfinyl, alkylsulfonyl, heteroarylthio, heteroarylsulfinyl, heteroarylsulfonyl, -PO(R^{13})(R^{14}), (where R^{13} and R^{14} are independently alkyl, aryl, alkoxy, aryloxy, heteroaryl, heteroarylalkyl, heteroaryloxy, heteroarylalkoxy, cycloheteroalkyl, cycloheteroalkylalkyl, cycloheteroalkoxy, or cycloheteroalkylalkoxy); R^1 can also be aminocarbonyl (where the amino may optionally be substituted with one or two aryl, alkyl or heteroaryl groups); cyano, 1,1-(alkoxyl or aryloxy)₂alkyl (where the two aryl or alkyl substituents can be independently defined, or linked to one another to form a ring, such as 1,3-dioxane or 1,3-dioxolane, connected to L^1 (or L^2 in the case of R^2) at the 2-position); 1,3-dioxane or 1,3-dioxolane connected to L^1 (or L^2 in the case of R^2) at the 4-position.

The R^1 group may have from one to four substituents, which can be any of the R^3 groups or R^1 groups, and any of the preferred R^1 substituents set out below.

- R^1 may be substituted with the following preferred substituents: alkylcarbonylamino, cycloalkylcarbonylamino, arylcarbonylamino, heteroarylcarbonylamino, alkoxycarbonylamino, aryloxycarbonylamino,
- 5 heteroaryloxylcarbonylamino, uriedo (where the uriedo nitrogens may be substituted with alkyl, aryl or heteroaryl), heterocyclylcarbonylamino (where the heterocycle is connected to the carbonyl group via a nitrogen or carbon atom), alkylsulfonylamino,
- 10 arylsulfonylamino, heteroarylsulfonylamino,



where J is: CHR^{23} , $\text{--}\overset{\text{O}}{\underset{\text{O}}{\text{C}}}\text{--}$, $\text{--}\underset{\text{R}^{24}}{\text{CH}}\text{--}\underset{\text{R}^{25}}{\text{CH}}\text{--}$ or $\text{--}\underset{\text{R}^{24}}{\text{C}}=\underset{\text{R}^{25}}{\text{C}}\text{--}$;

- 15 R^{23} , R^{24} and R^{25} are independently hydrogen, alkyl, alkenyl, alkynyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl, or cycloalkylalkyl;

- R^{20} , R^{21} , R^{22} are independently hydrogen, halo, alkyl, alkenyl, alkoxy, aryloxy, aryl, arylalkyl,
- 20 alkylmercapto, arylmercapto, cycloalkyl, cycloalkylalkyl, heteroaryl, heteroarylalkyl, hydroxy or haloalkyl; and these preferred substituents may either be directly attached to R^1 , or attached via an alkylene chain at an open position.

- 25 R^2 is the same or different from R^1 and is independently any of the groups set out for R^1 , H, polyhaloalkyl (such as CF_3CH_2 , $\text{CF}_3\text{CF}_2\text{CH}_2$ or CF_3) or cycloheteroalkyl, and may be substituted with one to four of any of the groups defined for R^3 , or any of the
- 30 substituents preferred for R^1 .


L^1 is a linking group containing from 1 to 10 carbons in a linear chain (including alkylene, alkenylene or alkynylene), which may contain, within the linking chain

any of the following: one or two alkenes, one or two alkynes, an oxygen, an amino group optionally substituted with alkyl or aryl, an oxo group; and may be substituted with one to five alkyl or halo groups (preferably F).

5 L^2 may be the same or different from L^1 and may independently be any of the L^1 groups set out above or a single bond.

R^3 , $R^{3'}$, R^4 and $R^{4'}$ may be the same or different and are independently selected from H, halogen, CF_3 , haloalkyl, hydroxy, alkoxy, alkyl, aryl, alkenyl, alkenyloxy, alkynyl, 10 alkynyloxy, alkanoyl, nitro, amino, thiol, alkylthio, alkylsulfinyl, alkylsulfonyl, carboxy, alkoxycarbonyl, aminocarbonyl, alkylcarbonyloxy, alkylcarbonylamino, cycloheteroalkyl, cycloheteroalkylalkyl, cyano, Ar, Ar-alkyl, ArO, Ar-amino, Ar-thio, Ar-sulfinyl, Ar-sulfonyl, 15 Ar-carbonyl, Ar-carbonyloxy or Ar-carbonylamino, wherein Ar is aryl or heteroaryl and Ar may optionally include 1, 2 or 3 additional rings fused to Ar;


R^{3a} and R^{3b} are the same or different and are 20 independently any of the R^3 groups except hydroxy, nitro, amino or thio;




are the same or different and independently represent a 5 25 or 6 membered heteroaryl ring which may contain 1, 2, 3 or 4 heteroatoms in the ring which are independently N, S or O; and including N-oxides.



X (in the fluorenyl type ring) is a bond, or is one of the following groups:

30





5



10 wherein

Y is O, N-R⁶ or S;

n' is 0, 1 or 2;

R⁶ is H, lower alkyl, aryl, -C(O)-R¹¹ or -C(O)-O-R¹¹;

15

R⁷ and R⁸ are the same or different and are independently H, alkyl, aryl, halogen, -O-R¹², orR⁷ and R⁸ together can be oxygen to form a ketone;R⁹, R¹⁰, R^{9'} and R^{10'} are the same or different and are independently H, lower alkyl, aryl or -O-R¹¹;

20

R^{9''} and R^{10''} are the same or different and are independently H, lower alkyl, aryl, halogen or -O-R¹¹;R¹¹ is alkyl or aryl;R¹² is H, alkyl or aryl.

25


The following provisos apply to formula I preferred compounds:

(a) when R¹ is unsubstituted alkyl or unsubstituted arylalkyl, L¹ cannot contain amino;30 (b) when R¹ is alkyl, L¹ cannot contain amino and oxo in adjacent positions (to form an amido group);(c) when R²L²A- is H₂N-, R¹L¹ cannot contain amino;(d) when R¹ is cyano, L¹ must have more than 2 carbons;(e) R¹L¹ must contain at least 3 carbons.


With respect to compounds IA and IB, R^2L^2 cannot have an O or N atom directly attached to $S=(O)_q$ or $CR^x(OH)$, and for IA, R^2L^2 cannot be H.

With respect to preferred compounds of formula IA and IB, where R^1 is cycloheteroalkyl, R^1 is exclusive of 1-piper-idinyl, 1-pyrrolidinyl, 1-azetidiny1 or 1-(2-oxo-pyrrolidinyl).

The MTP inhibitors disclosed in U.S. provisional application No. 60/017,253, filed May 10, 1996, (file HX82*) are pyrrolidine compounds and have the structure I



II



where Q is $-\overset{\text{O}}{\underset{\text{O}}{\text{C}}}-$ or $-\overset{\text{O}}{\underset{\text{O}}{\text{S}}}-$;

W is H, H or O;


X is: CHR^8 , $-\overset{\text{O}}{\underset{\text{O}}{\text{C}}}-$, $-\underset{\text{R}^9}{\text{CH}}-\underset{\text{R}^{10}}{\text{CH}}-$ or $-\underset{\text{R}^9}{\text{C}}=\underset{\text{R}^{10}}{\text{C}}-$;

R^8 , R^9 and R^{10} are independently hydrogen, alkyl, alkenyl, alkynyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl, or cycloalkylalkyl;


R^1 is alkyl, alkenyl, alkynyl, aryl, heteroaryl, arylalkyl (wherein alkyl preferably has at least 2 carbons, more preferably at least 3 carbons), diarylalkyl, arylalkenyl, diarylalkenyl, arylalkynyl, diarylalkynyl, diarylalkylaryl, heteroarylalkyl (wherein alkyl preferably has at least 2 carbons, more preferably at least 3 carbons), cycloalkyl, or cycloalkylalkyl (wherein alkyl preferably has at least 2 carbons, more preferably at least 3 carbons); all of the aforementioned R^1 groups being

optionally substituted through available carbon atoms with
 1, 2, 3 or 4 groups selected from halo, haloalkyl, alkyl,
 alkenyl, alkoxy, aryloxy, aryl, arylalkyl, alkylmercapto,
 arylmercapto, cycloalkyl, cycloalkylalkyl, heteroaryl,
 5 fluorenyl, heteroarylalkyl, hydroxy or oxo; or


R^1 is a fluorenyl-type group of the structure




10




R^1 is an indenyl-type group of the structure



15




Z^1 and Z^2 are the same or different and are independently a bond, O, S,




- 5 with the proviso that with respect to B , at least one of Z^1 and Z^2 will be other than a bond;

R^{11} is a bond, alkylene, alkenylene or alkynylene of up to 10 carbon atoms, arylene (for example



- 10 or mixed arylene-alkylene (for example



where n is 1 to 6;

R^{12} is hydrogen, alkyl, alkenyl, aryl, haloalkyl, trihaloalkyl, trihaloalkylalkyl, heteroaryl,

- 15 heteroarylalkyl, arylalkyl, arylalkenyl, cycloalkyl, aryloxy, alkoxy, arylalkoxy or cycloalkylalkyl; with the provisos that (1) when R^{12} is H, aryloxy, alkoxy or arylalkoxy, then Z^2 is
- $$-\text{NH}-\begin{array}{c} \text{C} \\ \parallel \\ \text{O} \end{array}, \quad \begin{array}{c} \text{N} \\ | \\ \text{alkyl} \end{array}-\begin{array}{c} \text{C} \\ \parallel \\ \text{O} \end{array}, \quad -\begin{array}{c} \text{C} \\ \parallel \\ \text{O} \end{array}- \quad \text{or a bond};$$


- 20 and (2) when Z^2 is a bond, R^{12} cannot be heteroaryl or heteroarylalkyl;

Z is a bond, O, S, N-alkyl, N-aryl, or alkylene or alkenylene of from 1 to 5 carbon atoms;


- 25 R^{13} , R^{14} , R^{15} , and R^{16} are independently hydrogen, alkyl, halo, haloalkyl, aryl, cycloalkyl, cycloheteroalkyl, alkenyl, alkynyl, hydroxy, alkoxy, nitro, amino, thio, alkylsulfonyl, arylsulfonyl, alkylthio, arylthio, aminocarbonyl, alkylcarbonyloxy, arylcarbonylamino, alkylcarbonylamino, arylalkyl, heteroaryl, heteroarylalkyl,
- 30 or aryloxy;

R^{15a} and R^{16a} are independently any of the R^{15} or R^{16} groups except hydroxy, nitro, amino or thio;

or R^1 is




wherein p is 1 to 8 and R¹⁷ and R¹⁸ are each independently H, alkyl, alkenyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl or cycloalkylalkyl, at least
 5 one of R¹⁷ and R¹⁸ being other than H;
 or R¹ is



wherein R¹⁹ is aryl or heteroaryl;
 R²⁰ is aryl or heteroaryl;
 10 R²¹ is H, alkyl, aryl, alkylaryl, arylalkyl, aryloxy, arylalkoxy, heteroaryl, heteroarylalkyl, heteroarylalkoxy, cycloalkyl, cycloalkylalkyl or cycloalkylalkoxy;
 R², R³, R⁴ are independently hydrogen, halo, alkyl,
 15 alkenyl, alkoxy, aryloxy, aryl, arylalkyl, alkylmercapto, arylmercapto, cycloalkyl, cycloalkylalkyl, heteroaryl, heteroarylalkyl, hydroxy or haloalkyl;
 R⁵ is alkyl, alkenyl, alkynyl, aryl, alkoxy, aryloxy, arylalkoxy, heteroaryl, arylalkyl,
 20 heteroarylalkyl, cycloalkyl, cycloheteroalkyl, heteroaryloxy, cycloalkylalkyl, polycycloalkyl, polycycloalkylalkyl, cycloalkenyl, cycloalkenylalkyl, polycycloalkenyl, polycycloalkenylalkyl, heteroarylcarbonyl, amino, alkylamino, arylamino,
 25 heteroarylamino, cycloalkyloxy, cycloalkylamino, all of the R⁵ substituents and R⁶ substituents (set out hereinafter) being optionally substituted through available carbon atoms with 1, 2, 3 or 4 groups selected from hydrogen, halo, alkyl, haloalkyl, alkoxy, haloalkoxy, alkenyl, alkynyl,
 30 cycloalkyl, cycloalkylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl, aryl, heteroaryl, arylalkyl, arylcycloalkyl, arylalkenyl, arylalkynyl, aryloxy, aryloxyalkyl, arylalkoxy, arylazo, heteroaryloxo, heteroarylalkyl, heteroarylalkenyl, heteroaryloxy, hydroxy,

nitro, cyano, amino, substituted amino (wherein the amino includes 1 or 2 substituents which are alkyl, aryl or heteroaryl, or any of the other aryl compounds mentioned in the definitions), thiol, alkylthio, arylthio, heteroarylthio, arylthioalkyl, alkylcarbonyl, arylcarbonyl, arylaminocarbonyl, alkoxycarbonyl, aminocarbonyl, alkynylaminocarbonyl, alkylaminocarbonyl, alkenylaminocarbonyl, alkylcarbonyloxy, arylcarbonyloxy, alkylcarbonylamino, arylcarbonylamino, arylsulfinyl, arylsulfinylalkyl, arylsulfonyl, alkylsulfonyl, arylsulfonylamino, heteroarylcarbonylamino, heteroarylsulfinyl, heteroarylthio, heteroarylsulfonyl, or alkylsulfinyl. Where R^5 is phenyl, aryl, heteroaryl or cycloalkyl; this group preferably includes an ortho hydrophobic substituent such as alkyl, haloalkyl (with up to 5 halo groups), alkoxy, haloalkoxy (with up to 5 halo groups), aryl, aryloxy or arylalkyl;

R^6 is hydrogen or C_1 - C_4 alkyl or C_1 - C_4 alkenyl;




are the same or different and are independently selected from heteroaryl containing 5- or 6-ring members; and including N-oxides of the formulae I and II compounds, that is




including pharmaceutically acceptable salts thereof.

The MTP inhibitors disclosed in U.S. provisional application No. 60/017,254, filed May 10, 1996, (file HX84*) are azetidine compounds which have the structure

I



II



5


where Q is $\text{—}\overset{\text{O}}{\underset{\text{O}}{\text{C}}}\text{—}$ or $\text{—}\overset{\text{O}}{\underset{\text{O}}{\text{S}}}\text{—}$;

X is: CHR^8 , $\text{—}\overset{\text{O}}{\underset{\text{O}}{\text{C}}}\text{—}$, $\text{—}\underset{\text{R}^9}{\text{CH}}\text{—}\underset{\text{R}^{10}}{\text{CH}}\text{—}$ or $\text{—}\underset{\text{R}^9}{\text{C}}=\underset{\text{R}^{10}}{\text{C}}\text{—}$;
n is 0 or 1;

10 R^8 , R^9 and R^{10} are independently hydrogen, alkyl, alkenyl, alkynyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl, or cycloalkylalkyl;



15 R^1 is alkyl, alkenyl, alkynyl, aryl, heteroaryl, arylalkyl (wherein alkyl preferably has at least 2 carbons, more preferably at least 3 carbons), diarylalkyl, arylalkenyl, diarylalkenyl, arylalkynyl, diarylalkynyl, diarylalkylaryl, heteroarylalkyl (wherein alkyl preferably has at least 2 carbons, more preferably at least 3 carbons), cycloalkyl, or cycloalkylalkyl (wherein alkyl preferably has at least 2 carbons, more preferably at least 3 carbons); all of the aforementioned R^1 groups being
20 optionally substituted through available carbon atoms with 1, 2, 3 or 4 groups selected from halo, haloalkyl, alkyl, alkenyl, alkoxy, aryloxy, aryl, arylalkyl, alkylmercapto, arylmercapto, cycloalkyl, cycloalkylalkyl, heteroaryl, fluorenyl, heteroarylalkyl, hydroxy or oxo; or
25

R^1 is a fluorenyl-type group of the structure




5

R^1 is an indenyl-type group of the structure




10

Z^1 and Z^2 are the same or different and are independently a bond, O, S,




15 with the proviso that with respect to B, at least one of Z^1 and Z^2 will be other than a bond;

R¹¹ is a bond, alkylene, alkenylene or alkynylene of up to 10 carbon atoms, arylene (for example



or mixed arylene-alkylene (for example



5

where q is 1 to 6;

R¹² is hydrogen, alkyl, alkenyl, aryl, haloalkyl, trihaloalkyl, trihaloalkylalkyl, heteroaryl, heteroarylalkyl, arylalkyl, arylalkenyl, cycloalkyl, aryloxy, alkoxy, arylalkoxy or cycloalkylalkyl; with the provisos that (1) when R¹² is H, aryloxy, alkoxy or arylalkoxy, then Z² is

$$\begin{array}{c} \text{—NH—C—} \\ \parallel \\ \text{O} \end{array}, \quad \begin{array}{c} \text{—N—C—} \\ \parallel \\ \text{alkyl O} \end{array}, \quad \begin{array}{c} \text{—C—} \\ \parallel \\ \text{O} \end{array} \quad \text{or a bond;}$$


and (2) when Z² is a bond, R¹² cannot be heteroaryl or heteroarylalkyl;

Z is a bond, O, S, N-alkyl, N-aryl, or alkylene or alkenylene of from 1 to 5 carbon atoms;

R¹³, R¹⁴, R¹⁵, and R¹⁶ are independently hydrogen, alkyl, halo, haloalkyl, aryl, cycloalkyl, cycloheteroalkyl, alkenyl, alkynyl, hydroxy, alkoxy, nitro, amino, thio, alkylsulfonyl, arylsulfonyl, alkylthio, arylthio, aminocarbonyl, alkylcarbonyloxy, arylcarbonylamino, alkylcarbonylamino, arylalkyl, heteroaryl, heteroarylalkyl, or aryloxy;


R^{15a} and R^{16a} are independently any of the R¹⁵ or R¹⁶ groups except hydroxy, nitro, amino or thio;

or R¹ is



wherein p is 1 to 8 and R¹⁷ and R¹⁸ are each independently H, alkyl, alkenyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl or cycloalkylalkyl, at least one of R¹⁷ and R¹⁸ being other than H;

or R¹ is



wherein R¹⁹ is aryl or heteroaryl;

R²⁰ is aryl or heteroaryl;


R²¹ is H, alkyl, aryl, alkylaryl, arylalkyl,

5 aryloxy, arylalkoxy, heteroaryl, heteroarylalkyl, heteroarylalkoxy, cycloalkyl, cycloalkylalkyl or cycloalkylalkoxy;


R², R³, R⁴ are independently hydrogen, halo, alkyl, alkenyl, alkoxy, aryloxy, aryl, arylalkyl, alkylmercapto, 10 arylmercapto, cycloalkyl, cycloalkylalkyl, heteroaryl, heteroarylalkyl, hydroxy or haloalkyl;

R⁵ is alkyl, alkenyl, alkynyl, aryl, alkoxy, aryloxy, arylalkoxy, heteroaryl, arylalkyl, heteroarylalkyl, cycloalkyl, cycloheteroalkyl, 15 heteroaryloxy, cycloalkylalkyl, polycycloalkyl, polycycloalkylalkyl, cycloalkenyl, cycloalkenylalkyl, polycycloalkenyl, polycycloalkenylalkyl, heteroarylcarbonyl, amino, alkylamino, arylamino, heteroarylamino, cycloalkyloxy, cycloalkylamino, all of the 20 R⁵ substituents and R⁶ substituents (set out hereinafter) being optionally substituted through available carbon atoms with 1, 2, 3 or 4 groups selected from hydrogen, halo, alkyl, haloalkyl, alkoxy, haloalkoxy, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, cycloheteroalkyl, 25 cycloheteroalkylalkyl, aryl, heteroaryl, arylalkyl, arylcycloalkyl, arylalkenyl, arylalkynyl, aryloxy, aryloxyalkyl, arylalkoxy, arylazo, heteroaryloxy, heteroarylalkyl, heteroarylalkenyl, heteroaryloxy, hydroxy, nitro, cyano, amino, substituted amino (wherein the amino 30 includes 1 or 2 substituents which are alkyl, aryl or heteroaryl, or any of the other aryl compounds mentioned in the definitions), thiol, alkylthio, arylthio, heteroarylthio, arylthioalkyl, alkylcarbonyl, arylcarbonyl, arylaminocarbonyl, alkoxycarbonyl, aminocarbonyl, 35 alkynylaminocarbonyl, alkylaminocarbonyl,


- alkenylaminocarbonyl, alkylcarbonyloxy, arylcarbonyloxy, alkylcarbonylamino, arylcarbonylamino, arylsulfinyl, arylsulfinylalkyl, arylsulfonyl, alkylsulfonyl, arylsulfonylamino, heteroarylcarbonylamino, heteroarylsulfinyl, heteroarylthio, heteroarylsulfonyl, or alkylsulfinyl. Where R^5 is phenyl, aryl, heteroaryl or cycloalkyl; this group preferably includes an ortho hydrophobic substituent such as alkyl, haloalkyl (with up to 5 halo groups), alkoxy, haloalkoxy (with up to 5 halo groups), aryl, aryloxy or arylalkyl;
- R^6 is hydrogen or C_1 - C_4 alkyl or C_1 - C_4 alkenyl;



- are the same or different and are independently selected from heteroaryl containing 5- or 6-ring members; and including N-oxides of the formulae I and II compounds, that is



- including pharmaceutically acceptable salts thereof.
- The MTP inhibitors disclosed in U.S. provisional application No. 60/028,216, filed October 1, 1996, have the structure




- including pharmaceutically acceptable salts thereof, wherein

A is (1) a bond;
(2) -O- ; or




where R^5 is H or lower alkyl or R^5 together with R^2 forms a carbocyclic or heterocyclic ring system containing 4 to 8 members in the ring.

5 B is a fluorenyl-type group of the structure:



10 B is an indenyl-type group of the structure



R^1 is independently alkyl, alkenyl, alkynyl, aryl, alkoxy, aryloxy, arylalkoxy, heteroaryl, arylalkyl, heteroarylalkyl, cycloalkyl, cycloalkylalkyl, polycycloalkyl, polycycloalkylalkyl, cycloalkenyl, cycloheteroalkyl, heteroaryloxy, cycloalkenylalkyl, polycycloalkenyl, polycycloalkenylalkyl, heteroarylcarbonyl, amino, alkylamino, arylamino,

15

20


heteroaryl-amino, cycloalkyloxy, cycloalkyl-amino, all optionally substituted through available carbon atoms with 1, 2, 3 or 4 groups selected from hydrogen, halo, alkyl, haloalkyl, alkoxy, haloalkoxy, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl, aryl, heteroaryl, arylalkyl, arylcycloalkyl, arylalkenyl, arylalkynyl, aryloxy, aryloxyalkyl, arylalkoxy, arylazo, heteroaryloxy, heteroarylalkyl, heteroarylalkenyl, heteroaryloxy, hydroxy, nitro, cyano, amino, substituted amino, thiol, alkylthio, arylthio, heteroarylthio, arylthioalkyl, alkylcarbonyl, arylcarbonyl, arylaminocarbonyl, alkoxycarbonyl, aminocarbonyl, alkynylaminocarbonyl, alkylaminocarbonyl, alkenylaminocarbonyl, alkylcarbonyloxy, arylcarbonyloxy, alkylcarbonylamino, arylcarbonylamino, arylsulfinyl, arylsulfinylalkyl, arylsulfonyl, alkylsulfonyl, arylsulfonylamino, heteroarylcarbonylamino, heteroarylsulfinyl, heteroarylthio, heteroarylsulfonyl, alkylsulfinyl;

20 R^2 is alkyl, alkenyl, alkynyl, alkoxyl, (alkyl or aryl)₃Si (where each alkyl or aryl group is independent), cycloalkyl, cycloalkenyl, substituted alkyl-amino, substituted arylalkyl-amino, aryl, arylalkyl, aryl-amino, aryloxy, heteroaryl, heteroaryl-amino, heteroaryloxy, arylsulfonylamino, heteroarylsulfonylamino, arylthio, arylsulfinyl, arylsulfonyl, alkylthio, alkylsulfinyl, alkylsulfonyl, heteroarylthio, heteroarylsulfinyl, heteroarylsulfonyl, -PO(R¹³)(R¹⁴), (where R¹³ and R¹⁴ are independently alkyl, aryl, alkoxy, aryloxy, heteroaryl, heteroarylalkyl, heteroaryloxy, heteroarylalkoxy, cycloheteroalkyl, cycloheteroalkylalkyl, cycloheteroalkoxy, or cycloheteroalkylalkoxy); R¹ can also be aminocarbonyl (where the amino may optionally be substituted with one or two aryl, alkyl or heteroaryl groups); cyano, 1,1-(alkoxyl or aryloxy)₂alkyl (where the two aryl or alkyl substituents can be independently defined, or linked to one another to

form a ring, such as 1,3-dioxane or 1,3-dioxolane, connected to L¹ (or L² in the case of R²) at the 2-position); 1,3-dioxane or 1,3-dioxolane connected to L¹ (or L² in the case of R²) at the 4-position.

5 The R² group may have from one to four substituents, which can be any of the R³ groups or R² groups, and any of the preferred R² substituents set out below.

R² may be substituted with the following preferred substituents: alkylcarbonylamino, cycloalkylcarbonylamino, 10 arylcarbonylamino, heteroarylcarbonylamino, alkoxycarbonylamino, aryloxycarbonylamino, heteroaryloxy carbonylamino, uriedo (where the uriedo nitrogens may be substituted with alkyl, aryl or heteroaryl), heterocyclylcarbonylamino (where the 15 heterocycle is connected to the carbonyl group via a nitrogen or carbon atom), alkylsulfonylamino, arylsulfonylamino, heteroarylsulfonylamino,



20

where J is: CHR^{23} , $\text{—}\overset{\text{O}}{\underset{\text{O}}{\text{C}}}\text{—}$, $\text{—}\underset{\text{R}^{24}}{\text{CH}}\text{—}\underset{\text{R}^{25}}{\text{CH}}\text{—}$ or $\text{—}\underset{\text{R}^{24}}{\text{C}}=\underset{\text{R}^{25}}{\text{C}}\text{—}$;

R²³, R²⁴ and R²⁵ are independently hydrogen, alkyl, alkenyl, alkynyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl, or cycloalkylalkyl;

25 R²⁰, R²¹, R²² are independently hydrogen, halo, alkyl, alkenyl, alkoxy, aryloxy, aryl, arylalkyl, alkylmercapto, arylmercapto, cycloalkyl, cycloalkylalkyl, heteroaryl, heteroarylalkyl, hydroxy or haloalkyl; and these preferred substituents may either be directly 30 attached to R¹, or attached via an alkylene chain at an open position.


L¹ is a linking group containing from 1 to 10 carbons in a linear chain (including alkylene, alkenylene or alkynylene), which may contain, within the linking chain

any of the following: one or two alkenes, one or two alkynes, an oxygen, an amino group optionally substituted with alkyl or aryl, an oxo group; and may be substituted with one to five alkyl or halo groups (preferably F).

5 L^2 may be the same or different from L^1 and may independently be any of the L^1 groups set out above or a single bond.

R^3 , $R^{3'}$, R^4 and $R^{4'}$ may be the same or different and are independently selected from H, halogen, CF_3 , haloalkyl, hydroxy, alkoxy, alkyl, aryl, alkenyl, alkenyloxy, alkynyl, 10 alkynyloxy, alkanoyl, nitro, amino, thiol, alkylthio, alkylsulfinyl, alkylsulfonyl, carboxy, alkoxycarbonyl, aminocarbonyl, alkylcarbonyloxy, alkylcarbonylamino, cycloheteroalkyl, cycloheteroalkylalkyl, cyano, Ar, 15 Ar-alkyl, ArO, Ar-amino, Ar-thio, Ar-sulfinyl, Ar-sulfonyl, Ar-carbonyl, Ar-carbonyloxy or Ar-carbonylamino, wherein Ar is aryl or heteroaryl and Ar may optionally include 1, 2 or 3 additional rings fused to Ar;


R^{3a} and R^{3b} are the same or different and are 20 independently any of the R^3 groups except hydroxy, nitro, amino or thio;



are the same or different and independently represent a 5 25 or 6 membered heteroaryl ring which may contain 1, 2, 3 or 4 heteroatoms in the ring which are independently N, S or O; and including N-oxides.



X (in the fluorenyl type ring) is a bond, or is one of the following groups:

30





5



10 wherein

Y is O, N-R⁶ or S;

n' is 0, 1 or 2;

R⁶ is H, lower alkyl, aryl, -C(O)-R¹¹ or -C(O)-O-R¹¹;15 R⁷ and R⁸ are the same or different and are independently H, alkyl, aryl, halogen, -O-R¹², orR⁷ and R⁸ together can be oxygen to form a ketone;R⁹, R¹⁰, R^{9'} and R^{10'} are the same or different and are independently H, lower alkyl, aryl or -O-R¹¹;20 R^{9''} and R^{10''} are the same or different and are independently H, lower alkyl, aryl, halogen or -O-R¹¹;R¹¹ is alkyl or aryl;R¹² is H, alkyl or aryl.

25 Compounds disclosed as preferred in each of the above applications are preferred for use in the present invention.



Most preferred MTP inhibitors to be employed in accordance with the present invention include preferred MTP inhibitors as set out in U.S. patent application Serial No. 548,811, filed January 11, 1996 (file DC21h) and in U.S. provisional application No. 60/017,224, filed May 9, 1996 (file HX79a*).

Thus, preferred compounds in U.S. patent application Serial No. 548,811 (file DC21h) for use herein are compounds



where Z is a bond;


5 X^1 and X^2 are H;


R^5 is aryl such as phenyl substituted with




(3) halo such as Cl

10 R^5 is heteroaryl such as  or  substituted with




wherein the R^5 substituent is preferably in the position adjacent to the carbon linked to .

15 $(CH_2)_x$ is $-(CH_2)_4-$ or




Most preferred is

9-[4-[4-[[2-(2,2,2-Trifluoroethoxy)benzoyl]amino]-1-piperidiny]butyl]-
N-(2,2,2-trifluoroethyl)-9H-fluorene-9-carboxamide




5

Preferred compounds in U.S. provisional application
No. 60/017,224 (file HX79a*) for use herein are MTP
inhibitor compounds of formula I that is



10 wherein A is NH,
B is



X is a bond, oxygen or sulfur; R³ and R⁴ are
independently H or F.

15 Preferred R¹ groups are aryl, preferably phenyl,
heteroaryl, preferably imidazolyl or pyridyl (preferably
substituted with one of the preferred R¹ substituents:
arylcarbonylamino, heteroarylcarbonylamino,
cycloalkylcarbonylamino, alkoxycarbonylamino,
20 alkylsulfonylamino, arylsulfonylamino,
heteroarylsulfonylamino), PO(Oalkyl)₂, heteroarylthio,
benzthi-azole-2-thio, imidazole-2-thio, alkyl, or alkenyl,
cycloalkyl such as cyclohexyl, or 1,3-dioxan-2-yl.

Preferred R² groups are alkyl, polyfluoroalkyl (such
25 as 1,1,1-trifluoroethyl), alkenyl, aryl or heteroaryl

(preferably substituted with one of the preferred R¹ substituents above), or PO(OAlkyl)₂.

If R² is alkyl, 1,1,1-trifluoroethyl, or alkenyl, it is preferred that R¹ is other than alkyl or alkenyl.

5 It is preferred that L¹ contains 1 to 5 atoms in the linear chain and L² is a bond or lower alkylene.

Preferred embodiments of formula IA and formula IB compounds of the invention include those where B, L¹, L², R¹ and R² are as set out with respect to the preferred
10 embodiments of the formula I compounds, q is 0 or 2 and R^x is H.


The other cholesterol lowering drugs or delipidating drugs which may be used in the method of the invention include HMG CoA reductase inhibitors, squalene synthetase
15 inhibitors, fibric acid derivatives, bile acid sequestrants, probucol, niacin, niacin derivatives and the like.


The HMG CoA reductase inhibitors suitable for use herein include, but are not limited to, mevastatin and
20 related compounds as disclosed in U.S. Patent No. 3,983,140, lovastatin (mevinolin) and related compounds as disclosed in U.S. Patent No. 4,231,938, pravastatin and related compounds such as disclosed in U.S. Patent No. 4,346,227, simvastatin and related compounds as disclosed
25 in U.S. Patent Nos. 4,448,784 and 4,450,171, with pravastatin, lovastatin or simvastatin being preferred. Other HMG CoA reductase inhibitors which may be employed herein include, but are not limited to, fluvastatin, cerivastatin, atorvastatin, pyrazole analogs of
30 mevalonolactone derivatives as disclosed in U.S. Patent No. 4,613,610, indene analogs of mevalonolactone derivatives as disclosed in PCT application WO 86/03488, 6-[2-(substituted-pyrrol-1-yl)alkyl]pyran-2-ones and derivatives thereof as disclosed in U.S. Patent No.
35 4,647,576, Searle's SC-45355 (a 3-substituted pentanedioic acid derivative) dichloroacetate, imidazole analogs of mevalonolactone as disclosed in PCT application WO

86/07054, 3-carboxy-2-hydroxy-propane-phosphonic acid derivatives as disclosed in French Patent No. 2,596,393, 2,3-di-substituted pyrrole, furan and thiophene derivatives as disclosed in European Patent Application No. 0221025, 5 naphthyl analogs of mevalonolactone as disclosed in U.S. Patent No. 4,686,237, octahydronaphthalenes such as disclosed in U.S. Patent No. 4,499,289, keto analogs of mevinolin (lovastatin) as disclosed in European Patent Application No. 0,142,146 A2, as well as other known HMG 10 CoA reductase inhibitors.

In addition, phosphinic acid compounds useful in inhibiting HMG CoA reductase suitable for use herein are disclosed in GB 2205837.

The squalene synthetase inhibitors suitable for use 15 herein include, but are not limited to, α -phosphonosulfonates disclosed in U.S. application Serial No. 08/266,888, filed July 5, 1994 (HX59b), those disclosed by Biller et al, J. Med. Chem. 1988, Vol. 31, No. 10, pp 1869-1871, including isoprenoid 20 (phosphinylmethyl)phosphonates such as those of the formula



R¹

5

including the triacids thereof, triesters thereof and tripotassium and trisodium salts thereof as well as other squalene synthetase inhibitors disclosed in U.S. Patent
 10 Nos. 4,871,721 and 4,924,024 and in Biller et al, J. Med. Chem., 1988, Vol. 31, No. 10, pp 1869 to 1871.

In addition, other squalene synthetase inhibitors suitable for use herein include the terpenoid pyrophosphates disclosed by P. Ortiz de Montellano et al,
 15 J. Med. Chem.; 1977, 20, 243-249, the farnesyl diphosphate analog A and presqualene pyrophosphate (PSQ-PP) analogs as disclosed by Corey and Volante, J. Am. Chem. Soc. 1976, 98, 1291-1293, phosphinylphosphonates reported by McClard, R.W. et al, J.A.C.S., 1987, 109, 5544 and cyclopropanes reported
 20 by Capson, T.L., PhD dissertation, June, 1987, Dept. Med. Chem. U. of Utah, Abstract, Table of Contents, pp. 16, 17, 40-43, 48-51, Summary.

Preferred are pravastatin, lovastatin or simvastatin.

25 All of the above U.S. applications are incorporated herein by reference.

Other cholesterol lowering drugs suitable for use herein include, but are not limited to, antihyperlipoproteinemic agents such as fibric acid derivatives, such as fenofibrate, gemfibrozil, clofibrate, bezafibrate, ciprofibrate, clinofibrate and the like, probucol, and related compounds as disclosed in U.S. Patent No. 3,674,836, probucol and gemfibrozil being preferred, bile acid sequestrants such as cholestyramine, colestipol and DEAE-Sephadex (Secholex®, Polidexide®), as well as clofibrate, lipostabil (Rhone-Poulenc), Eisai E-5050 (an N-substituted ethanolamine derivative), imanixil (HOE-402), tetrahydrolipstatin (THL), istigmastanylphosphorylcholine (SPC, Roche), aminocyclodextrin (Tanabe Seiyoku), Ajinomoto AJ-814 (azulene derivative), melinamide (Sumitomo), Sandoz 58-035, American Cyanamid CL-277,082 and CL-283,546 (disubstituted urea derivatives), nicotinic acid, acipimox, acifran, neomycin, p-aminosalicylic acid, aspirin, poly(diallylmethylamine) derivatives such as disclosed in U.S. Patent No. 4,759,923, quaternary amine poly(diallyldimethylammonium chloride) and ionenes such as disclosed in U.S. Patent No. 4,027,009, and other known serum cholesterol lowering agents.

The above-mentioned U.S. patents are incorporated herein by reference.

The MTP inhibitor will be employed in a weight ratio to the other cholesterol lowering or delipidating agent (where present), in accordance with the present invention, within the range from about 500:1 to about 1:500, preferably from about 100:1 to about 1:100.

In carrying out the method of the present invention, the MTP inhibitor alone or optionally in combination with the other cholesterol lowering drug may be administered to mammalian species, such as monkeys, dogs, cats, rats, humans, etc., and as such may be incorporated in a conventional systemic dosage form, such as a tablet, capsule, elixir or injectable. The above dosage forms will also include the necessary carrier material, excipient,

lubricant, buffer, antibacterial, bulking agent (such as mannitol), anti-oxidants (ascorbic acid or sodium bisulfite) or the like. Oral dosage forms are preferred, although parenteral forms are quite satisfactory as well.

5 The dose administered must be carefully adjusted according to age, weight and condition of the patient, as well as the route of administration, dosage form and regimen and the desired result.

10 The dosages and formulations for the MTP inhibitor will be as disclosed in the various patents and applications discussed above.

15 The dosages and formulations for the other delipidating agent to be employed, where applicable, will be as set out in the latest edition of the Physicians' Desk Reference.

For oral administration, a satisfactory result may be obtained employing the MTP inhibitor in an amount within the range of from about 0.01 mg/kg to about 100 mg/kg and preferably from about 0.1 mg/kg to about 75 mg/kg, one to 20 four times daily.

A preferred oral dosage form, such as tablets or capsules, will contain the MTP inhibitor in an amount of from about 5 to about 500 mg, preferably from about 10 to about 400 mg, and more preferably from about 20 to about 25 250 mg, one to four times daily.

For parenteral administration, the MTP inhibitor will be employed in an amount within the range of from about 0.005 mg/kg to about 10 mg/kg and preferably from about 0.005 mg/kg to about 8 mg/kg, one to four times 30 daily.

The other cholesterol lowering or delipidating agent will be employed in amounts set out in the latest edition of the Physician's Desk Reference (PDR).

35 For oral administration, a satisfactory result may be obtained employing an HMG CoA reductase inhibitor in dosages employed, for example, pravastatin, lovastatin,

simvastatin, atorvastatin, fluvastatin or cerivastatin as indicated in the Physician's Desk Reference, such as in an amount within the range of from about 1 to 2000 mg, and preferably from about 4 to about 200 mg.

5 The squalene synthetase inhibitor may be employed in dosages in an amount within the range of from about 10 mg to about 2000 mg and preferably from about 25 mg to about 200 mg.

10 A preferred oral dosage form, such as tablets or capsules, will contain the HMG CoA reductase inhibitor in an amount of from about 0.1 to about 100 mg, preferably from about 5 to about 80 mg, and more preferably from about 10 to about 40 mg.

15 A preferred oral dosage form, such as tablets or capsules will contain the squalene synthetase inhibitor in an amount of from about 10 to about 500 mg, preferably from about 25 to about 200 mg.

20 The serum cholesterol lowering drugs when present will be employed in dosages normally employed as indicated in the Physician's Desk Reference, for each of such agents such as in an amount within the range of from about 2 mg to about 7500 mg and preferably from about 2 mg to about 4000 mg.

25 A preferred oral dosage form, such as tablets or capsules, will contain MTP inhibitor in an amount of from about 10 to about 400 mg, and the HMG CoA reductase inhibitor (where present) in an amount of from about 0.1 to about 100 mg, preferably from about 5 to about 80 mg, and more preferably from about 10 to about 50 mg.

30 The MTP inhibitor and other cholesterol lowering drug may be employed together in the same oral dosage form or in separate oral dosage forms taken at the same time.

35 The compositions described above may be administered in the dosage forms as described above in single or divided doses of one to four times daily. It may be advisable to start a patient on a low dose combination and work up gradually to a high dose combination.

Tablets of various sizes can be prepared, e.g., of about 2 to 2000 mg in total weight, containing one or both of the active substances in the ranges described above, with the remainder being a physiologically acceptable carrier of other materials according to accepted pharmaceutical practice. These tablets can, of course, be scored to provide for fractional doses. Gelatin capsules can be similarly formulated.

Liquid formulations can also be prepared by dissolving or suspending one or the combination of active substances in a conventional liquid vehicle acceptable for pharmaceutical administration so as to provide the desired dosage in one to four teaspoonsful.

Such dosage forms can be administered to the patient on a regimen of one to four doses per day.

According to another modification, in order to more finely regulate the dosage schedule, the active substances may be administered separately in individual dosage units at the same time or carefully coordinated times. Since blood levels are built up and maintained by a regulated schedule of administration, the same result is achieved by the simultaneous presence of the two substances. The respective substances can be individually formulated in separate unit dosage forms in a manner similar to that described above.

Fixed combinations of MTP inhibitor and other cholesterol lowering drug are more convenient and are preferred, especially in tablet or capsule form for oral administration.

In formulating the compositions, the active substances, in the amounts described above, are compounded according to accepted pharmaceutical practice with a physiologically acceptable vehicle, carrier, excipient, binder, preservative, stabilizer, flavor, etc., in the particular type of unit dosage form.

Illustrative of the adjuvants which may be incorporated in tablets are the following: a binder such as

gum tragacanth, acacia, corn starch or gelatin; an excipient such as dicalcium phosphate or cellulose; a disintegrating agent such as corn starch, potato starch, alginic acid or the like; a lubricant such as stearic acid or magnesium stearate; a sweetening agent such as sucrose, aspartame, lactose or saccharin; a flavoring agent such as orange, peppermint, oil of wintergreen or cherry. When the dosage unit form is a capsule, it may contain in addition to materials of the above type a liquid carrier such as a fatty oil. Various other materials may be present as coatings or to otherwise modify the physical form of the dosage unit. For instance, tablets or capsules may be coated with shellac, sugar or both. A syrup or elixir may contain the carrier, glycerol as solubilizer, sucrose as sweetening agent, methyl and propyl parabens as preservatives, a dye and a flavoring such as cherry or orange.

Some of the active substances described above form commonly known, pharmaceutically acceptable salts such as alkali metal and other common basic salts or acid addition salts, etc. References to the base substances are therefore intended to include those common salts known to be substantially equivalent to the parent compound.

The formulations as described above will be administered for a prolonged period, that is, for as long as phytosterolemia exists. Sustained release forms of such formulations which may provide such amounts biweekly, weekly, monthly and the like may also be employed.

The following Examples represent preferred embodiments of the present invention.

Examples 1 and 2

Formulations suitable for oral administration are prepared as described below.

Capsules each containing about 5 mg MTP inhibitor
 5 BMS 201,038 (Example 1) and capsules each containing about 50 mg BMS 201,038 (Example 2) are produced from the following ingredients.

10	<u>Ingredient</u>	<u>Example 1</u>		<u>Example 2</u>	
		<u>Amount (mg/ Capsule)</u>		<u>Amount (mg/ Capsule)</u>	
	BMS-201038 (1)	1.1		56.9	
	Lactose, Hydrous, NF ca.	30.2	ca.	99.9	
	Lactose, Anhydrous, NF	47.3		0.0	
	Microcrystalline Cellulose, NF	10.0		50.0	
	Pregelatinized Starch, NF	5.0		25.0	
	Sodium Starch Glycolate, NF	5.0		12.5	
	Colloidal Silicon Dioxide, NF	1.0		5.0	
	Magnesium Stearate, NF	0.3		0.6	
	Purified Water, USP or	q.s.		q.s.	
	Water for Injection, USP	q.s.		q.s.	
	Gray, Opaque, Size #0 Capsule Shell	One Capsule		One Capsule	
		about		about	
	Total Fill Weight	1000.0		250.0	

(1) In Example 1, this amount is expressed in terms of the amount of methane sulfonic acid salt per capsule at
 15 100% potency. In Example 2, this amount is expressed in terms of the free base. This is equivalent to 1 mg and 50 mg (Examples 1 and 2, respectively) of the free base.

The MTP inhibitor BMS 201,038, and colloidal silicon
 20 dioxide are blended in a suitable blender with lactose hydrous, microcrystalline cellulose, pregelatinized starch

and a portion of sodium starch glycolate. The resulting blend is wet granulated with water. The wet granulation is dried in a suitable dryer. The remaining portion of sodium starch glycolate is added to the granulation and mixed
5 therein. Magnesium stearate is added to the granulation and mixed therein. The resulting blend is filled into capsules.

Example 3

10 Pravastatin tablets (10, 20 or 40 mg as described in the 1996 PDR) and MTP inhibitor (BMS 201,238) tablets may be administered as a combination in accordance with the teachings of the present invention. In addition, the pravastatin and MTP inhibitor tablets may be ground up into
15 powders and used together in a single capsule.

Example 4

Tablets containing 500 mg clofibrate by itself or in combination with 10 mg BMS 201,038 may be employed in
20 separate dosage forms or combined in a single capsule form.

Examples 5, 6 and 7


Ciprofibrate, bezafibrate, gemfibrozil alone or in combination with an MTP inhibitor may also be prepared in a
25 manner described hereinbefore in Examples 1 to 3.

What is claimed is:

1. A method for inhibiting onset of or treating
phytosterolemia in a mammalian species, which comprises
administering to a patient in need of treatment a
5 therapeutically effective amount of an MTP inhibitor.

2. The method as defined in Claim 1 wherein the MTP
inhibitor is employed in combination with another type of
cholesterol lowering drug.

3. The method as defined in Claim 2 wherein the MTP
10 inhibitor has the structure



including the piperidine N-oxide thereof or a
pharmaceutically acceptable salt thereof, wherein Z is a
15 bond, O or S;

X¹ and X² are independently selected from H or halo;

x is an integer from 2 to 6;

R⁵ is heteroaryl, aryl, heterocycloalkyl or
cycloalkyl, each R⁵ group being optionally substituted with
20 1, 2, 3 or 4 substituents which may be the same or
different.

4. The method as defined in Claim 3 where in the
MTP inhibitor Z is a bond.

5. The method as defined in Claim 3 where the MTP
25 inhibitor is a piperidine N-oxide.

6. The method as defined in Claim 3 where in the
MTP inhibitor (CH₂)_x is optionally substituted with 1, 2 or
3 substituents which are the same or different and are
alkyl or halo.




7. The method as defined in Claim 3 where in the
30 MTP inhibitor R⁵ is substituted with 1, 2, 3 or 4
substituents which may be the same or different and are

halogen, monocyclic heteroaryl, bicyclic heteroaryl,
 heteroarylalkyl, cycloheteroalkyl, alkyl, alkoxy,
 cycloalkyl, aryl, aryloxy, substituted aryl, arylalkyloxy,
 heteroaryloxy, amino, alkylamino, alkyl(aryl)amino,
 5 heteroarylamino, arylamino, alkylthio, arylthio,
 arylthioalkyl, heteroarylthio, arylsulfinyl or acyl.


8. The method as defined in Claim 7 where in the
 MTP inhibitor the R⁵ includes a substituent attached to a
 carbon in the position adjacent to the carbon linked to $\overset{\text{O}}{\parallel}{\text{C}}$.


10 9. The method as defined in Claim 7 where in the
 MTP inhibitor R⁵ is substituted with 1, 2, 3 or 4 of one or
 more of the following


I, Cl, F, CF₃








alkyl, phenyl, phenyl substituted with halo, alkyl, CF₃O,



alkoxy, , CF₃, or phenyl; $\text{---}\overset{\text{H}}{\text{N}}\text{---}(\text{CH}_2)_p\text{CF}_3\text{---}$ where p is 1 to 5, -N(CH₃)C₆H₅;

-S-(CH₂)_pCF₃ where p is 1 to 5, , -S-alkyl,






5 $\text{---}\text{S---}(\text{CH}_2)_p\text{---}\overset{\text{O}}{\parallel}\text{S---}\overset{\text{O}}{\parallel}\text{---C}_6\text{H}_5$, , OCH₃;

cyclohexyl, , ;

amino, , , ;


, ;

10 alkanoyl, alkoxycarbonyl, aroyl, heteroarylaminocarbonyl, arylalkyloxycarbonyl,


, $\text{---}\text{N}(\text{CH}_3)\text{---CH}_2\text{---}$ , $\text{---CH}_2\text{---S---C}_6\text{H}_5$, $\text{---O---CH}_2\text{---}$ , ---S--- , ---O--- , $\text{---S---}\overset{\text{O}}{\parallel}\text{---C}_6\text{H}_4\text{---Cl}$.

15 10. The method as defined in Claim 9 where in the MTP inhibitor R⁵ is phenyl substituted with haloalkylphenyl or heteroaryl.


11. The method as defined in Claim 10 where in the MTP inhibitor R⁵ is



12. The method as defined in Claim 9 where in the MTP inhibitor is




or



5

13. The method as defined in Claim 2 wherein the MTP inhibitor has the structure



10 including pharmaceutically acceptable salts thereof,
N-oxides thereof,

wherein q is 0, 1 or 2;

A is (1) a bond;

(2) -O-; or



(3) $\begin{array}{c} \text{---N---} \\ | \\ \text{R}^5 \end{array}$

15



where R^5 is H or lower alkyl, or R^5 together with R^2 forms a carbocyclic or heterocyclic ring system containing 4 to 8 members in the ring;

B is a fluorenyl-type group of the structure

20




5 B is an indenyl-type group of the structure



- 10 R^x is H, alkyl or aryl;
 R^1 is alkyl, alkenyl, alkynyl, alkoxy, (alkyl or aryl)₃Si (where each alkyl or aryl group is independent), cycloalkyl, cycloalkenyl, substituted alkylamino, substituted arylalkylamino, aryl, arylalkyl, arylamino,
 15 aryloxy, heteroaryl, heteroarylamino, heteroaryloxy, arylsulfonylamino, heteroarylsulfonylamino, arylthio, arylsulfinyl, arylsulfonyl, alkylthio, alkylsulfinyl, alkylsulfonyl, heteroarylthio, heteroarylsulfinyl, heteroarylsulfonyl, $-PO(R^{13})(R^{14})$, (where R^{13} and R^{14} are
 20 independently alkyl, aryl, alkoxy, aryloxy, heteroaryl, heteroarylalkyl, heteroaryloxy, heteroarylalkoxy, cycloheteroalkyl, cycloheteroalkylalkyl, cycloheteroalkoxy,

or cycloheteroalkylalkoxy); aminocarbonyl (where the amino may optionally be substituted with one or two aryl, alkyl or heteroaryl groups); cyano, 1,1-(alkoxyl or aryloxy)₂alkyl (where the two aryl or alkyl substituents
 5 can be independently defined, or linked to one another to form a ring connected to L¹ (or L² in the case of R²) at the 2-position); 1,3-dioxane or 1,3-dioxolane connected to L¹ (or L² in the case of R²) at the 4-position; the R¹ group may optionally be substituted with 1, 2, 3 or 4
 10 substituents, which can be any of the R³ or R¹ groups or alkylcarbonylamino, cycloalkylcarbonylamino, arylcarbonylamino, heteroarylcarbonylamino, alkoxycarbonylamino, aryloxycarbonylamino, heteroaryloxycarbonylamino, uriedo (where the uriedo
 15 nitrogens may optionally be substituted with alkyl, aryl or heteroaryl), heterocyclylcarbonylamino (where the heterocycle is connected to the carbonyl group via a nitrogen or carbon atom), alkylsulfonylamino, arylsulfonylamino, heteroarylsulfonylamino,

20



where J is: CHR²³, $\begin{array}{c} \text{---C---} \\ || \\ \text{O} \end{array}$, $\begin{array}{c} \text{---CH---CH---} \\ | \quad | \\ \text{R}^{24} \quad \text{R}^{25} \end{array}$ or $\begin{array}{c} \text{---C=C---} \\ | \quad | \\ \text{R}^{24} \quad \text{R}^{25} \end{array}$;

R²³, R²⁴ and R²⁵ are independently hydrogen, alkyl, alkenyl,
 25 alkynyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl, or cycloalkylalkyl;

R²⁰, R²¹, R²² are independently hydrogen, halo, alkyl, alkenyl, alkoxy, aryloxy, aryl, arylalkyl, alkylmercapto, arylmercapto, cycloalkyl, cycloalkylalkyl,
 30 heteroaryl, heteroarylalkyl, hydroxy or haloalkyl; and these substituents may either be directly attached to R¹, or attached via an alkylene at an open position;

R² is independently any of the groups set out for R¹, H, polyhaloalkyl, or cycloheteroalkyl, and may be


optionally substituted with one to four of any of the groups defined for R^3 or substituents defined for R^1 ;

L^1 is a linking group containing from 1 to 10 carbons in a linear chain including alkylene, alkenylene or alkynylene, which may contain, within the linking chain any of the following: one or two alkenes, one or two alkynes, an oxygen, an amino group, an oxo group, and may be substituted with one to five alkyl or halo groups;

L^2 may be the same or different from L^1 and may independently be any of the L^1 groups set out above or a single bond;


R^3 , $R^{3'}$, R^4 and $R^{4'}$ may be the same or different and are independently selected from H, halogen, CF_3 , haloalkyl, hydroxy, alkoxy, alkyl, aryl, alkenyl, alkenyloxy, alkynyl, alkynyloxy, alkanoyl, nitro, amino, thiol, alkylthio, alkylsulfinyl, alkylsulfonyl, carboxy, alkoxycarbonyl, aminocarbonyl, alkylcarbonyloxy, alkylcarbonylamino, cycloheteroalkyl, cycloheteroalkylalkyl, cyano, Ar-, Ar-alkyl, ArO, Ar-amino, Ar-thio, Ar-sulfinyl, Ar-sulfonyl, Ar-carbonyl, Ar-carbonyloxy or Ar-carbonylamino, wherein Ar is aryl or heteroaryl and Ar may optionally include 1, 2 or 3 additional rings fused to Ar;

R^{3a} and R^{3b} are the same or different and are independently any of the R^3 groups except hydroxy, nitro, amino or thio;





are the same or different and independently represent a 5 or 6 membered heteroaryl ring which contains 1, 2, 3 or 4 heteroatoms in the ring which are independently N, S or O; and including N-oxides;

X is a bond, or is one of the following groups:







5



10



wherein

Y is O, N-R⁶ or S;

n' is 0, 1 or 2;

15 R⁶ is H, lower alkyl, aryl, -C(O)-R¹¹ or -C(O)-O-R¹¹;

R⁷ and R⁸ are the same or different and are independently H, alkyl, aryl, halogen, -O-R¹², or

R⁷ and R⁸ together can be oxygen to form a ketone;

20 R⁹, R¹⁰, R^{9'} and R^{10'} are the same or different and are independently H, lower alkyl, aryl or -O-R¹¹;

R^{9''} and R^{10''} are the same or different and are independently H, lower alkyl, aryl, halogen or -O-R¹¹;

25 R¹¹ is alkyl or aryl;

R¹² is H, alkyl or aryl;

14. The method as defined in Claim 13 wherein the



30 15. The method as defined in Claim 14 wherein A is a bond.


16. The method as defined in Claim 14 wherein A is -O-.

17. The method as defined in Claim 14 wherein A is




18. The method as defined in Claim 14 wherein B is a fluorenyl-type group.

19. The method as defined in Claim 14 having the formula




wherein B is



A is NH;
X is a bond, oxygen or sulfur;
R³ and R⁴ are the same or different and are H or F;
R¹ is aryl, phenyl, heteroaryl, imidazolyl, pyridyl, cyclohexyl, PO(R¹³)(R¹⁴), heteroarylthio, benzthiazole-2-thio, imidazole-2-thio, alkyl, alkenyl or 1,3-dioxan-2-yl, wherein each of the above is optionally substituted;
R² is alkyl, polyfluoroalkyl, alkenyl, aryl, phenyl, heteroaryl, imidazolyl or pyridyl, wherein each of the above is optionally substituted;
L¹ is a chain containing 1 to 5 atoms in a linear chain;
L² is a bond or lower alkylene.

20. The method as defined in Claim 1 wherein the MTP inhibitor has the structure



including pharmaceutically acceptable salts thereof, N-oxides thereof,

wherein



A is (1) a bond;

(2) $-O-$; or

(3) $\begin{array}{c} \text{---N---} \\ | \\ R^5 \end{array}$



5 where R^5 is H or lower alkyl, or R^5 together with R^2 forms a carbocyclic or heterocyclic ring system containing 4 to 8 members in the ring;

B is a fluorenyl-type group of the structure



B is an indenyl-type group of the structure

15




20

R^x is H, alkyl or aryl;

R^1 is independently alkyl, alkenyl, alkynyl, aryl, alkoxy, aryloxy, arylalkoxy, heteroaryl, arylalkyl, heteroarylalkyl, cycloalkyl, cycloalkylalkyl, polycycloalkyl, polycycloalkylalkyl, cycloalkenyl, cycloheteroalkyl, heteroaryloxy, cycloalkenylalkyl, polycycloalkenyl, polycycloalkenylalkyl, heteroarylcarbonyl, amino, alkylamino, arylamino, heteroarylamino, cycloalkyloxy, cycloalkylamino, all optionally substituted through available carbon atoms with
 10 1, 2, 3 or 4 groups selected from hydrogen, halo, alkyl, haloalkyl, alkoxy, haloalkoxy, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl, aryl, heteroaryl, arylalkyl, arylcycloalkyl, arylalkenyl, arylalkynyl, aryloxy, aryloxyalkyl, arylalkoxy, arylazo, heteroaryloxy, heteroarylalkyl, heteroarylalkenyl, heteroaryloxy, hydroxy, nitro, cyano, amino, substituted amino, thiol, alkylthio, arylthio, heteroarylthio, arylthioalkyl, alkylcarbonyl, arylcarbonyl, arylaminocarbonyl, alkoxycarbonyl, aminocarbonyl, alkynylaminocarbonyl, alkylaminocarbonyl, alkenylaminocarbonyl, alkylcarbonyloxy, arylcarbonyloxy, alkylcarbonyl-amino, arylcarbonylamino, arylsulfinyl, arylsulfinylalkyl, arylsulfonyl, alkylsulfonyl, arylsulfonylamino, heteroarylcarbonylamino,
 20 heteroarylsulfinyl, heteroarylthio, heteroarylsulfonyl, alkylsulfinyl;

R^2 is alkyl, alkenyl, alkynyl, alkoxyl, (alkyl or aryl)₃Si (where each alkyl or aryl group is independent), cycloalkyl, cycloalkenyl, substituted alkylamino, substituted arylalkylamino, aryl, arylalkyl, arylamino, aryloxy, heteroaryl, heteroarylamino, heteroaryloxy, arylsulfonylamino, heteroarylsulfonylamino, arylthio, arylsulfinyl, arylsulfonyl, alkylthio, alkylsulfinyl, alkylsulfonyl, heteroarylthio, heteroarylsulfinyl, heteroarylsulfonyl, -PO(R^{13})(R^{14}), (where R^{13} and R^{14} are
 35 independently alkyl, aryl, alkoxy, aryloxy, heteroaryl, heteroarylalkyl, heteroaryloxy, heteroarylalkoxy,

- cycloheteroalkyl, cycloheteroalkylalkyl, cycloheteroalkoxy, or cycloheteroalkylalkoxy); aminocarbonyl (where the amino may optionally be substituted with one or two aryl, alkyl or heteroaryl groups); cyano, 1,1-(alkoxyl or aryloxy)₂alkyl (where the two aryl or alkyl substituents can be independently defined, or linked to one another to form a ring connected to L¹ (or L² in the case of R²) at the 2-position); 1,3-dioxane or 1,3-dioxolane connected to L¹ (or L² in the case of R²) at the 4-position; the R² group may optionally be substituted with 1, 2, 3 or 4 substituents, which can be any of the R³ or R¹ groups or alkylcarbonylamino, cycloalkylcarbonylamino, arylcarbonylamino, heteroarylcarbonylamino, alkoxycarbonylamino, aryloxycarbonylamino, heteroaryloxylcarbonylamino, uriedo (where the uriedo nitrogens may optionally be substituted with alkyl, aryl or heteroaryl), heterocyclylcarbonylamino (where the heterocycle is connected to the carbonyl group via a nitrogen or carbon atom), alkylsulfonylamino, arylsulfonylamino, heteroarylsulfonylamino,



where J is: CHR^{23} , $\begin{array}{c} \text{—C—} \\ \parallel \\ \text{O} \end{array}$, $\begin{array}{c} \text{—CH—CH—} \\ | \quad | \\ \text{R}^{24} \text{ R}^{25} \end{array}$ or $\begin{array}{c} \text{—C=C—} \\ | \quad | \\ \text{R}^{24} \text{ R}^{25} \end{array}$;


- R²³, R²⁴ and R²⁵ are independently hydrogen, alkyl, alkenyl, alkynyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloalkyl, or cycloalkylalkyl;
R²⁰, R²¹, R²² are independently hydrogen, halo, alkyl, alkenyl, alkoxy, aryloxy, aryl, arylalkyl, alkylmercapto, arylmercapto, cycloalkyl, cycloalkylalkyl, heteroaryl, heteroarylalkyl, hydroxy or haloalkyl; and these substituents may either be directly attached to R¹, or attached via an alkylene at an open position;

L¹ is a linking group containing from 1 to 10 carbons in a linear chain including alkylene, alkenylene or alkynylene, which may contain, within the linking chain any of the following: one or two alkenes, one or two alkynes, an oxygen, an amino group, an oxo group, and may be substituted with one to five alkyl or halo groups;

L² may be the same or different from L¹ and may independently be any of the L¹ groups set out above or a single bond;


R³, R^{3'}, R⁴ and R^{4'} may be the same or different and are independently selected from H, halogen, CF₃, haloalkyl, hydroxy, alkoxy, alkyl, aryl, alkenyl, alkenyloxy, alkynyl, alkynyloxy, alkanoyl, nitro, amino, thiol, alkylthio, alkylsulfinyl, alkylsulfonyl, carboxy, alkoxycarbonyl, aminocarbonyl, alkylcarbonyloxy, alkylcarbonylamino, cycloheteroalkyl, cycloheteroalkylalkyl, cyano, Ar-, Ar-alkyl, ArO, Ar-amino, Ar-thio, Ar-sulfinyl, Ar-sulfonyl, Ar-carbonyl, Ar-carbonyloxy or Ar-carbonylamino, wherein Ar is aryl or heteroaryl and Ar may optionally include 1, 2 or 3 additional rings fused to Ar;


R^{3a} and R^{3b} are the same or different and are independently any of the R³ groups except hydroxy, nitro, amino or thio;





are the same or different and independently represent a 5 or 6 membered heteroaryl ring which contains 1, 2, 3 or 4 heteroatoms in the ring which are independently N, S or O; and including N-oxides;

X is a bond, or is one of the following groups:





5



10 wherein

Y is O, N-R⁶ or S;

n' is 0, 1 or 2;

R⁶ is H, lower alkyl, aryl, -C(O)-R¹¹ or -C(O)-O-R¹¹;

15

R⁷ and R⁸ are the same or different and are independently H, alkyl, aryl, halogen, -O-R¹², orR⁷ and R⁸ together can be oxygen to form a ketone;R⁹, R¹⁰, R^{9'} and R^{10'} are the same or different and are independently H, lower alkyl, aryl or -O-R¹¹;

20

R^{9''} and R^{10''} are the same or different and are independently H, lower alkyl, aryl, halogen or -O-R¹¹;R¹¹ is alkyl or aryl;R¹² is H, alkyl or aryl.

25

21. The method as defined in Claim 2 wherein the other cholesterol lowering drug is an HMG CoA reductase inhibitor, a squalene synthetase inhibitor, a fibric acid derivative, probucol, a bile acid sequestrant, nicotinic acid or neomycin.

30

22. The method as defined in Claim 21 wherein the HMG CoA reductase inhibitor is pravastatin, lovastatin, simvastatin, atorvastatin, fluvastatin or cerivastatin.

23. The method as defined in Claim 2 wherein the cholesterol lowering drug is a fibric acid derivative which is gemfibrozil, fenofibrate, clofibrate, bezafibrate, ciprofibrate, clinofibrate, probucol, gemfibrozil,
5 dextrothyroxine or its sodium salt, colestipol or its hydrochloride, cholestyramine, nicotinic acid, neomycin, p-aminosalicylic acid or aspirin.

24. The method as defined in claim 2 wherein the MTP inhibitor is present in a weight ratio to the other
10 cholesterol lowering drug (where present) within the range from about 500:1 to about 1:500.

25. The method as defined in Claim 1 wherein the MTP inhibitor lowers plasma LDL-cholesterol to at least 50% of normal LDL blood levels, and lowers triglycerides to at
15 least 50% of normal triglyceride blood levels.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/00618

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A01N 43/40; A61K 31/445

US CL : 514/324, 325, 326, 321, 320, 319

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/324, 325, 326, 321, 320, 319

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
NONE

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y,E	US 5,712,279 A (BILLER et al.) 27 January 1998, see abstract	1-25
A	US 5,502,045 A (MIETTINEN et al.) 26 March 1996, see abstract.	1-25
Y	Database SCISEARCH on STN, COBB et al., Abstract to, "Sitosterolemia-opposing effects of cholestyramine and lovastatin on plasma sterol levels in a homozygous girl and her heterozygous father", (Rockefeller Univ., Biochem. Genet & Metab Lab, New York, NY), Abstract No. 96:434984, Metabolism-Clinical and Experimental, Vol. 45, No. 6, pp. 673-679, June 1996.	1-25

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

08 APRIL 1998

Date of mailing of the international search report

07 MAY 1998

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized Officer

DWAYNE C. JONES

Telephone No. (703) 308-1235

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/00618

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

REGISTRY, CAPLUS, USPATFULL, BIOSIS, MEDLINE, EMBASE, LIFESCI, BIOTECHDS, WPIDS, CONFSCI, DISSABS, SCISEARCH, JICST-EPLUS, PROMT, TOXLIT, TOXLINE, DRUGU, DRUGNL, DRUGLAUNCH

search terms: structure search with the following terms, (mtp or protein)(w)(inhibitor or piperidine(w)oxide, inhibit? or prevent? or treat? or therap?